

Bend Magnet Heat Loads & Out of Orbit Scenarios

Timothy Valicenti¹, Jason Carter², Kamlesh Suthar^{2,*}

1. Brown University: Dept. of Mech. Engineering,
2. Argonne National Laboratory: AES
*Project Supervisor


Challenge

- Accelerated electrons emit radiation.
- This radiation can heat up materials.
- True electron paths contain errors.
- How do these errors effect the power distribution?

Tasks

- Calculate the ideal path an electron travels through a given bend magnet.
- Determine sets of alternative paths the electron may take due to orbital errors.
- Create ray traces of photons emitted from the electrons.
- Calculate the power distributions on the surfaces impacted by the photons.
- Verify data with SynRad simulations

Process




Accomplishments

- Task 1: Solved for any ideal path
- Task 2: Code finds desired off orbit paths
- Task 3: Can ray trace from any trajectory
- Task 4: Calculates heat map on any plane
- Task 5: Data closely matches SynRad


Future Work

- A more friendly UI
- Deeper COMSOL Integration
- More geometries of absorbers
- Insertion device analogs

Geometry




Theory



$$\mathcal{A}_u = \gamma_x x^2 + \alpha_x x x' + \beta_x x'^2$$


$$\mathcal{A}_u = \gamma_y y^2 + \alpha_y y y' + \beta_y y'^2$$

$$\begin{bmatrix} z(t) \\ x(t) \\ y(t) \end{bmatrix} = \bar{R}(\theta) \begin{bmatrix} \rho \sin(\omega t) \\ \rho(\cos(\omega t) - 1) \\ 0 \end{bmatrix} + \bar{r}_0$$






$$\frac{\partial^2 P}{\partial \theta \partial \psi} = P_0 \frac{1}{(1 + X^2)^{5/2}} \left[1 + \frac{5}{7} \frac{X^2}{1 + X^2} \right], \quad X = \gamma \theta$$

$$P_0 \left(\frac{W}{\text{mrad}^2} \right) = 5.421 * E^4 (\text{GeV}) * I(A) * B(T)$$



Performance & Discussion

- Data matches SynRad to high degrees of accuracy and yet doesn't contain the noise.
- Can be used to determine what the heat load is on an absorbing surface of a specific material.
- One can run an optimization process by changing the absorber position and orientation to reduce the peak intensity.
- Analytical solutions allow easy coupling with other programs such as COMSOL.



Acknowledgements

A very special thanks to Kamlesh Suthar for his guidance on this project; to Jason Carter, Jason Lerch, Kathy Harkay, and Roger Dejus for their insight along the way; and especially to Pat Den Hartog, Eric Prebys, and Linda Spentzouris for giving me this opportunity and organizing the Summer 2016 student Internships.

References

- Capatina, Dana. private communication (2016).
- Carter, Jason. private communication (2016).
- Chao, Alexander Wu., and M. Tigner. *Handbook of Accelerator Physics and Engineering*. River Edge, NJ: World Scientific, 1999. Print.
- Dejus, Roger. "Power Distribution from a Dipole Source." *Internal APS Memo* (2003): 1-8. Print.
- Edwards, D. A., and M. J. Syphers. *An Introduction to the Physics of High Energy Accelerators*. New York: Wiley, 1993. Print.
- Harkay, Katherine. "Maximum Beam Orbit in MBA and Ray Tracing Guidelines." 2nd ser. (2014): 1-9. Print.
- Suthar, Kamlesh. private communication (2016).